


#0009: MRT-6160, a VAV1-directed molecular glue degrader, attenuates T and B cell effector functions and inhibits disease progression in a spontaneous autoimmune MRL-Fas^{lpr} mouse model

Cartwright ANR², Gyger L², Desai F¹, Vora S¹, Kostikova A², Wang X¹, Trenh P¹, May K¹, Nguyen S¹, Chris King¹, Lam D¹, Lucas X², Zlotosch M¹, Liardo E², Lamberto I¹, Demarco B¹, Bonenfant D², Townson S¹, Krishnan E¹, Janku F¹, Castle J¹, McAllister L², Paterson A¹, Peluso M¹

¹Monte Rosa Therapeutics Inc., 321 Harrison Ave, Boston, MA 02118, United States

²Monte Rosa Therapeutics AG, WKL-136.3, Klybeckstrasse 191, 4057 Basel, Switzerland

VAV1 is a guanine nucleotide exchange factor with a critical role in T- and B-cell receptor signaling and activity




MRT-6160 is a rationally designed molecular glue degrader that selectively degrades VAV1 in human and mouse immune cells

Molecular glue degraders (MGD) function to induce structural changes in ubiquitin ligases, such as cereblon, to drive the formation of ternary structures with a target protein.


Following binding of cereblon to the target, this protein is then ubiquitin tagged and subsequently degraded via the proteasome-mediated degradation machinery of the cell.

MGDs can induce degradation of otherwise 'undruggable' proteins as the mechanism does not require a classical binding pocket, contrary to conventional protein inhibitors, significantly increasing the target space and potential utility across a range of diseases.




Human PBMCs and mouse splenocytes were treated overnight with dose-range of MRT-6160, after which VAV1 protein levels were assessed by JESS. Percentage (%) VAV1 degradation was calculated by normalizing VAV1 expression to β -actin loading control and shown as relative to DMSO control. Data from N = 3 biological replicates. Human PBMCs and mouse splenocytes were treated for 24 hrs with 10 μ M MRT-6160 then assessed by quantitative tandem mass tag proteomics. The y-axis represents p-value [-log₁₀]; the x-axis represents protein fold change [log₂] relative to DMSO (0.1%) control samples. Dark blue circles represent CRBN neosubstrates including the target, VAV1, and other known cereblon neosubstrates; GSPT1, IKZF1, IKZF3, CSNK1A1 (CK1 α), SALL4, and ZFP91 or VAV family members; VAV2 and VAV3.

MRT-6160 degrades VAV1 and attenuates T and B cell effector functions in healthy and rheumatic disease patient donors




MRT-6160 decreases Tfh cell-mediated B cell activation, differentiation, and immunoglobulin secretion



Summary and Future Development

- MRT-6160 is a first-in-class VAV1 MGD that **attenuates TCR/BCR activity *in vitro* & *in vivo*** attenuating T/B cell effector functions and differentiation.
- Oral administration of MRT-6160 **attenuates proteinuria, lymphadenopathy, skin lesion formation, autoantibody production, organomegaly, and histopathology** in an MRL-Fas^{lpr} mouse model.
- Given its *in vitro* and *in vivo* MOA profile, MRT-6160 has **strong potential to alleviate disease symptoms in multiple autoimmune and inflammatory diseases** including rheumatic diseases such as Sjögren's disease, SLE, rheumatoid arthritis and others.
- MRT-6160 is exclusively licenced by Novartis and has **completed assessment in a Phase I healthy volunteer study (NCT06597799)**.

MRT-6160 inhibits key disease readouts in a spontaneous autoimmune MRL-Fas^{lpr} mouse model



MRT-6160 reduces kidney glomerular and interstitial nephritis in MRL-Fas^{lpr} mice

